



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Features of the mcr-1 cassette related to colistin resistance

Zurfluh, Katrin ; Kieffer, Nicolas ; Poirel, Laurent ; Nordmann, Patrice ; Stephan, Roger

DOI: <https://doi.org/10.1128/AAC.01519-16>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-129983>

Journal Article

Accepted Version

Originally published at:

Zurfluh, Katrin; Kieffer, Nicolas; Poirel, Laurent; Nordmann, Patrice; Stephan, Roger (2016). Features of the mcr-1 cassette related to colistin resistance. *Antimicrobial Agents and Chemotherapy*, 60:6438-6439.

DOI: <https://doi.org/10.1128/AAC.01519-16>

Antimicrobial Agents and Chemotherapy – New Data Letter

Features of the *mcr-1* cassette respect to colistin resistance

Katrin Zurfluh¹, Nicolas Kieffer², Laurent Poirel^{2,3}, Patrice Nordmann^{2,3,4}, Roger Stephan^{1#}

¹Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Switzerland.

²Emerging Antibiotic Resistance Unit, Medical and Molecular Microbiology, Department of Medicine, University of Fribourg, Fribourg, Switzerland, ³INSERM European Unit (LEA Paris, France), University of Fribourg, Fribourg, Switzerland, ⁴University of Lausanne and University Hospital Center, Lausanne, Switzerland.

[#]Corresponding author:

Roger Stephan, Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zurich, Winterthurerstr. 272, CH-8057 Zurich, Switzerland, Phone 0041-44-6358651, Fax 0041-44-6358908, e-mail stephanr@fsafety.uzh.ch

The recent description of the plasmid-mediated colistin resistance gene, *mcr-I*, in strains isolated from food animals, food and humans in China was a signal for an avalanche of retrospective and prospective studies investigating the occurrence of this specific gene (1). The *mcr-I* gene has been identified almost all over the world now, and the earliest evidence for its presence dates back to the 1980's (2). The *mcr-I* gene has so far been associated with non-related types of plasmid replicons such as IncI2, IncHI2, IncP, IncFIB and IncX4 (1,3-5) and was found only rarely to be chromosomally encoded (6). This gene is part of a 2,600-bp long fragment designed as the *mcr-I* cassette that encompasses the likely promoter sequences for *mcr-I* expression (7). The *mcr-I* gene is most often located at the right-hand extremity of the insertion element IS*AplI*, together with a 723-bp long open reading frame (named *orf723*) encoding an hypothetical protein. According to blast analysis, it is a putative phosphoesterase, and shares 45% and 44% identities with those of *Corynebacterium durum* (Accession-Nr: WP_060996190.1) and *Psychrobacter arcticus* (Accession-Nr: WP_011280438.1), respectively. However, the putative contribution of this orf for expression of the *mcr-I* gene and subsequently to colistin resistance remains unknown.

Our goal was to evaluate the role of *orf723* with respect to the colistin resistance. Therefore, three *Escherichia coli* recombinant strains were constructed, with the same plasmid harbouring either the *mcr-I* gene alone, *orf723* alone, and the entire *mcr-I* cassette, respectively. The primers used to amplify the *mcr-I* gene, the 723-bp orf and the whole *mcr-I* cassette are listed in Table 1, and the *mcr-I*-positive *E. coli* OW3E1 (GenBank Accession number: KX129783) was used as template. Amplicons were double-digested with restriction enzymes BamHI and EcoRI and cloned into the low-

copy vector pCCR9 (Taxonomy ID: 125570) digested with the respective enzymes, to create vectors pCCR9::mcr-1, pCCR9::orf723 and pCCR9::mcr-1::orf723. The constructs were transformed by electrotransformation into *E. coli* DH5 α , giving rise to recombinant strains DH5 α ::pCCR9::mcr-1, DH5 α ::pCCR9::orf723, and DH5 α ::pCCR9::mcr-1::orf723, respectively. Minimal inhibitory concentrations (MIC) of colistin were determined using broth dilution tests as recommended by EUCAST. MICs are summarized in Table 2. MIC values of recombinant strains expressing MCR-1 with and without the *orf723* were increased and identical. These results further confirm that expression of the *mcr-1* gene confers reduced susceptibility to colistin. However they show that *orf723* encoding an hypothetical protein and which has likely been co-mobilized with the *mcr-1* gene from its original genetic context does not impact colistin susceptibility.

Acknowledgments:

This work was supported by funding from the Universities of Zurich and Fribourg, Switzerland, and by the Swiss Federal Office of Public Health.

Transparency declarations

None to declare

References

1. **Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J.** 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: A microbiological and molecular biological study. *Lancet Infect. Dis.* **16**, 161–8.
2. **Schwarz S, Johnson AP.** 2016. Transferable resistance to colistin: a new but old threat. *J Antimicrob. Chemother.* **71**, 2066-70.
3. **Nordmann P, Lienhard R, Kieffer N, Clerc O, Poirel L.** 2016. Plasmid-mediated colistin-resistant *Escherichia coli* in bacteremia in Switzerland. *Clin. Infect. Dis.* **5**, 1322-3.
4. **Zhi C, Lv L, Yu LF, Doi Y, Liu JH.** 2016. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect. Dis.* **16**, 292-3.
5. **Zurfluh K, Klumpp, J, Nüesch-Inderbinen M, Stephan, R.** Full-length nucleotide sequences of *mcr-1* harboring plasmids isolated from extended-spectrum β -lactamase (ESBL)- producing *Escherichia coli* of different origins. *Antimicrob. Agents. Chemother.*, accepted.
6. **Zurfluh K, Tasara T, Poirel L, Nordmann, P, Stephan, R.** 2016. Draft genome sequence of *Escherichia coli* S51, a chicken isolate harboring a chromosomally-encoded *mcr-1* gene. *Genome Announc.* **4**:e00796-16; doi:10.1128/genomeA.00796-16.

89 7. **Poirel L, Kieffer N, Brink A, Coetze J, Jayol A, Nordmann P.** 2016. Genetic
90 features of MCR-1-producing colistin-resistant *Escherichia coli* isolates in South
91 Africa. *Antimicrob. Agents Chemother.* **60**, 4394-7.

92

93

Table 1: Primers used for cloning of *mcr-I*, the hypothetical protein and the whole *mcr-I* cassette

Primer name	Sequence (5'-3') ^a	T _m (°C) ^b	Location
mcr_BamHI_up	TTTTTTGGATCCGCCGCAATTATCCCACCG	53	22 bp upstream of <i>mcr-I</i> start codon ^c
mcr_EcoRI_dn	TTTTTTGAATTCCCACCGCCCATACGAATGG	56	36 bp downstream of <i>mcr-I</i> stop codon
orf723_BamHI_up2	TTTTTTGGATCCGCACACTCCATTCGTATTATGGGC	57	18 bp upstream of 723 bp orf start codon
orf723_EcoRI_dn	TTTTTTGAATTCCCGTTCCTATTGGTAGTTTCCAGG	56	81 bp downstream of 723 bp orf stop codon

^a The restriction sites are underlined.

^b T_m, melting point.

^c downstream of putative promotor region.

123

124 Table 2: Minimal inhibitory concentrations (MIC's) of colistin for each *E. coli* DH5 α
125 conjugants as well as the negative controls *E. coli* DH5 α ::pCCR9 and *E. coli* DH5 α
126 using broth dilution tests as recommended by EUCAST
127
128

Conjugant / Strain	MIC colistin [mg/L]
DH5 α ::pCCR9::mcr-1	4
DH5 α ::pCCR9::orf723	1
DH5 α ::pCCR9::mcr-1::orf723	4
DH5 α ::pCCR9	0.5
DH5 α	1

129

130

131

132

133

134